

## ESSENTIAL PROPERTIES RELATED TO SHORT-TIME FRACTIONAL FOURIER TRANSFORM

SRI SULASTERI<sup>a\*</sup>, NASRULLAH BACHTIAR<sup>b</sup>, WAHYUNI EKASASMITA<sup>c</sup>

<sup>a</sup> *Department of Mathematics Education, Universitas Islam Negeri Alauddin Makassar,*

<sup>b</sup> *Department of Mathematics, Hasanuddin University,*

<sup>c</sup> *Department of Mathematics, Institut Teknologi Bacharuddin Jusuf Habibie,  
email : sri.sulasteri@uin-alauddin.ac.id, nasrullahmipa.013@gmail.com,  
wahyuni.ekasasmita@ith.ac.id*

Accepted June 24, 2024    Revised September 17, 2024    Published October 31, 2024

**Abstract.** *We start by defining the short-time fractional Fourier transform in this paper, which is a natural generalization of the fractional Fourier transform. We then investigate its essential properties and explore an uncertainty principle related to this proposed transformation.*

**Keywords:** Fractional Fourier transform, short-time fractional Fourier transform, uncertainty principle

### 1. Introduction

It is well known that the fractional Fourier transform (FrFT) is a generalized of the usual Fourier transform (FT), has several applications in optics, signal processing, and quantum mechanics [1,2,3,4,5,6,7]. The fractional Fourier transform depends on a parameter  $\vartheta$  and can be considered as a counterclockwise rotation of the function to arbitrary angles in the spatial frequency domain. Since the FrFT is an extension of the FT, we may transfer some properties of the FT to the FrFT domain like shifting, scaling, modulation, convolution, and inequalities with some changes [8,9,10].

On the other hand, the fractional Fourier domain-frequency contents, which are needed in some applications, are not found by FrFt. This problem can be solved using short-time fractional Fourier transform (STFrFT) [11]. A crucial timefrequency analysis tool is the STFrFT [12]. It has been extensively applied in numerous signal processing domains including speech and acoustics.

The uncertainty principle (UP) is a key concept in signal processing and quantum mechanics. In many changes, one of the well-liked outcomes is likewise the uncertainty principle. A number of works, including [13,14], have recently addressed

\*Corresponding author

the uncertainty principles associated with the conventional short-time Fourier transform. The STFrFT, which may be thought of as one rotation of the conventional short-time Fourier transform, was proposed by the authors in [15]. The authors of [16] conclude that there is a connection between the FrFT parameter and the uncertainty principle for the STFrFT by reducing a fundamental inequality. Nevertheless, there are currently no publications that include the crucial characteristics and underlying principles of uncertainty for the short-time fractional Fourier transform (STFrFT).

This paper’s remaining sections are arranged as follows. The fundamentals of the fractional Fourier transform (FrFT) are presented in Section 2, along with a summary of several results that are crucial to understanding the conclusions drawn in this work. The definition of the short-time fractional Fourier transform (STFrFT) is presented in Section 3. This section also includes the derivation of certain fundamental properties, such as Moyal’s and the inversion formula. The attributes will yield significant benefits for the advancement of our primary findings. We also investigate a principle of uncertainty associated with this suggested transformation. Lastly, we present our conclusion in Section 4.

**2. Basic Facts on Fractional Fourier Transform**

In this part, we mainly review the basic notations on the fractional Fourier transform (FrFT). We start with the well known definition below.

For  $1 \leq p \leq \infty$ , we use the notation  $L^s(\mathbb{R})$  to represent the space of measurable functions on real numbers  $\mathbb{R}$  with norm:

$$\|h\|_{L^s(\mathbb{R})} = \left( \int_{\mathbb{R}} |h(x)|^s dt \right)^{1/s} < \infty, \quad 1 \leq s < \infty. \tag{2.1}$$

In particular, when  $s = \infty$  we obtain  $L^\infty(\mathbb{R})$ -norm:

$$\|h\|_{L^\infty(\mathbb{R})} = \text{ess sup}_{x \in \mathbb{R}} |h(x)|. \tag{2.2}$$

Especially, if  $h \in L^\infty(\mathbb{R})$  is continuous then:

$$\|h\|_{L^\infty(\mathbb{R})} = \sup_{x \in \mathbb{R}} |h(x)|. \tag{2.3}$$

The usual inner product of  $L^2(\mathbb{R})$  is then defined as:

$$\langle h, \phi \rangle_{L^2(\mathbb{R})} = \int_{\mathbb{R}} h(x) \overline{\phi(x)} dx. \tag{2.4}$$

where  $\overline{\phi(x)}$  is complex conjugate of  $\phi(x)$ .

We remind that the Fourier transform of a function  $h \in L^2(\mathbb{R})$  is defined by:

$$\mathcal{F}\{h\}(\psi) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} h(x) e^{-ix\psi} dx. \tag{2.5}$$

**Definition 2.1.** For any function  $h \in L^2(\mathbb{R})$ , the fractional Fourier transform (FrFT) with parameter  $\vartheta$  is described by the formula

$$\mathcal{F}_\vartheta\{h\}(\psi) = \int_{\mathbb{R}} h(x) K_\vartheta(\psi, x) dx, \tag{2.6}$$

where the kernel transform  $K_{\vartheta}(\psi, x)$  is defined by:

$$K_{\vartheta}(\psi, x) = \begin{cases} C_{\vartheta} e^{i(|x|^2+|\psi|^2)\frac{\cot \vartheta}{2} - it \cdot \psi \csc \vartheta}, & \vartheta \neq n\pi, \\ \delta(x - \psi), & \vartheta = 2n\pi, \\ \delta(x + \psi), & \vartheta = (2n + 1)\pi, n \in \mathbb{Z}, \end{cases} \quad (2.7)$$

for  $\delta$  is Dirac delta function. Here

$$C_{\vartheta} = \sqrt{\frac{1 - i \cot \vartheta}{2\pi}}, \quad \overline{C_{\vartheta}} = \sqrt{\frac{1 + i \cot \vartheta}{2\pi}}, \quad (2.8)$$

and it is evident that:

$$|C_{\vartheta}| = \frac{1}{\sqrt{2\pi \sin \vartheta}}. \quad (2.9)$$

It is straightforward to verify that the FrFT kernel fulfills the following basic properties.

$$\overline{K_{\vartheta}(\psi, x)} = K_{-\vartheta}(\psi, x), \quad (2.10)$$

and

$$\int_{\mathbb{R}} K_{\vartheta}(\psi, x) \overline{K_{\vartheta}(\psi', t)} dt = \delta(\psi - \psi'), \quad (2.11)$$

where  $\overline{K_{\vartheta}(\psi, x)}$  is the complex conjugate of  $K_{\vartheta}(\psi, x)$ .

**Definition 2.2.** For any function  $h \in L^2(\mathbb{R})$  with  $\mathcal{F}_{\vartheta}\{h\} \in L^2(\mathbb{R})$ . The inversion formula of the fractional Fourier transform (FrFT) of the function  $h$  is described as:

$$\begin{aligned} h(x) &= \mathcal{F}_{\vartheta}^{-1}[\mathcal{F}_{\vartheta}\{h\}](x), \\ &= \int_{\mathbb{R}} \mathcal{F}_{\vartheta}\{h\}(\psi) \overline{K_{\vartheta}(\psi, x)} d\psi, \\ &= \int_{\mathbb{R}} \mathcal{F}_{\vartheta}\{h\}(\psi) \overline{C_{\vartheta}} e^{-i(|x|^2+|\psi|^2)\frac{\cot \vartheta}{2} + ix \cdot \psi \csc \vartheta} d\psi. \end{aligned} \quad (2.12)$$

Equation (2.12) tell us to recover the original function from its FrFT.

**Lemma 2.3 (FrFT Parseval [17]).** For two functions  $h, \phi \in L^2(\mathbb{R})$ , it holds

$$\langle h, \phi \rangle_{L^2(\mathbb{R})} = \langle \mathcal{F}_{\vartheta}\{h\}, \mathcal{F}_{\vartheta}\{\phi\} \rangle_{L^2(\mathbb{R})}, \quad (2.13)$$

and

$$\|h\|_{L^2(\mathbb{R})}^2 = \|\mathcal{F}_{\vartheta}\{h\}\|_{L^2(\mathbb{R})}^2. \quad (2.14)$$

### 3. Short-Time Fractional Fourier Transform and Its Essential Properties

This part begins by defining the short-time fractional Fourier transform (STFrFT) which is a generalized of the classical fractional Fourier transform. We then collect its important properties which will be needed later.

**Definition 3.1.** For any function  $\phi \in L^\infty(\mathbb{R})$  and  $h \in L^2(\mathbb{R})$ , the short-time fractional Fourier transform (STFrFT) with respect to  $\phi$  is described by

$$\begin{aligned} \mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) &= C_\vartheta \int_{\mathbb{R}} h(x) \overline{\phi(x-t)} e^{i\left(\frac{x^2+\psi^2}{2}\right) \cot \vartheta - ix\psi \csc \vartheta} dx \\ &= \int_{\mathbb{R}} h(x) \overline{\phi(x-t)} K_\vartheta(x, \psi) dx. \end{aligned} \quad (3.1)$$

For a fixed  $t$ , equation (3.1) may be expressed in the form

$$\mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) = \mathcal{F}_\vartheta\{\check{A}_{f,\bar{\phi}}\}(\psi), \quad (3.2)$$

where

$$\check{A}_{f,\bar{\phi}}(x, t) = h(x) \overline{\phi(x-t)}. \quad (3.3)$$

**Remark 3.2.** It should be observed that when  $\vartheta = \frac{\pi}{2}$ , equation (3.1) becomes:

$$\begin{aligned} \mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \check{A}_{h,\phi}(x, t) e^{-ix\psi} dx \\ &= \mathcal{F}\{\check{A}_{h,\phi}\}(\psi). \end{aligned} \quad (3.4)$$

The fundamental relationship of the STFrFT with classical FT is given by:

$$\begin{aligned} C_\vartheta^{-1} \mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) e^{-i\psi^2 \frac{\cot \vartheta}{2}} &= \int_{\mathbb{R}} \check{A}_{h,\bar{\phi}}^\vartheta(x, t) e^{-ix\psi \csc \vartheta} dx \\ &= \mathcal{F}\left\{\check{A}_{h,\bar{\phi}}^\vartheta(x, t)\right\}(\psi \csc \vartheta), \end{aligned} \quad (3.5)$$

where

$$\check{A}_{h,\bar{\phi}}^\vartheta(x, t) = h(x) \phi(x-t) e^{i\frac{t^2 \cot \vartheta}{2}}. \quad (3.6)$$

Some of the essential properties of the STFrFT are collected in the following theorems.

**Theorem 3.3 (Moyal's formula).** Let  $h, \phi \in L^2(\mathbb{R})$ , we have:

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) \right|^2 d\psi dt = \|h\|_{L^2(\mathbb{R})}^2 \|\phi\|_{L^2(\mathbb{R})}^2. \quad (3.7)$$

**Proof.** In view of Parseval's formula for the FrFT in equation (2.14) and relation (3.4), we have:

$$\begin{aligned} \int_{\mathbb{R}} \left| \mathcal{STF}_\phi^\vartheta\{h\}(t, \psi) \right|^2 d\psi &= \int_{\mathbb{R}} \left| \mathcal{F}\{\check{A}_{h,\phi}\}(\psi) \right|^2 d\psi, \\ &= \int_{\mathbb{R}} \left| \check{A}_{h,\phi}(\psi) \right|^2 dx, \\ &= \int_{\mathbb{R}} \check{A}_{h,\bar{\phi}}(\psi) \overline{\check{A}_{h,\bar{\phi}}(\psi)} d\psi, \\ &= \int_{\mathbb{R}} h(x) \overline{\phi(x-t)} \overline{h(x) \phi(x-t)} dx, \\ &= \int_{\mathbb{R}} h(x) \overline{h(x) \phi(x-t)} \phi(x-t) dx. \end{aligned} \quad (3.8)$$

We integrate the both sides of equataion (3.8) with respect to  $dt$  and get:

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right|^2 d\psi dt, = \int_{\mathbb{R}} \int_{\mathbb{R}} h(x) \overline{h(x)\phi(x-t)} \phi(x-t) dx dt. \tag{3.9}$$

Performing the change of variables  $x - t = u$ , we infer that:

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right|^2 d\psi dt = \int_{\mathbb{R}} \int_{\mathbb{R}} h(x) \overline{h(x)\phi(u)} \phi(u) dx du. \tag{3.10}$$

Using Fubini's theorem on the right side of equation (3.10), we further get:

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right|^2 d\psi dt = \int_{\mathbb{R}} |h(x)|^2 dx \int_{\mathbb{R}} |\phi(u)|^2 du, \tag{3.11}$$

or,

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) \right|^2 d\psi dt = \|h\|_{L^2(\mathbb{R})}^2 \|\phi\|_{L^2(\mathbb{R})}^2, \tag{3.12}$$

which completes the proof. □

**Theorem 3.4 (Inversion formula).** *Let  $h, \phi \in L^2(\mathbb{R})$ , then inversion formula of the STFrFT is given by:*

$$h(x) = \frac{1}{\phi(0)} \int_{\mathbb{R}} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) K_{-\vartheta}(x, \psi) dx. \tag{3.13}$$

**Proof.** It follows with relation (3.1) that:

$$\begin{aligned} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) &= C_{\vartheta} \int_{\mathbb{R}} h(x) \overline{\phi(x-t)} e^{i\left(\left(\frac{x^2+\psi^2}{2}\right) \cot \vartheta - ix\psi \csc \vartheta\right)} dx \\ &= \int_{\mathbb{R}} \check{A}_{f, \overline{\phi}}(x, t) K_{\vartheta}(x, \psi) dx, \end{aligned} \tag{3.14}$$

where  $\check{A}_{f, \overline{\phi}}(x, t)$  is defined by equation (3.3). Using inversion of FrFT in (2.12), then:

$$\check{A}_{f, \overline{\phi}}(x, t) = \int_{\mathbb{R}} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) K_{-\vartheta}(x, \psi) dx. \tag{3.15}$$

Equation equation (3.15) above can be rewritten in the form:

$$h(x) \overline{\phi(x-t)} = \int_{\mathbb{R}} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) K_{-\vartheta}(x, \psi) dx. \tag{3.16}$$

Choosing  $x = t$  in equation (3.16), we obtain

$$h(x) \overline{\phi(0)} = \int_{\mathbb{R}} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) K_{-\vartheta}(x, \psi) dx. \tag{3.17}$$

It means that

$$h(x) = \frac{1}{\phi(0)} \int_{\mathbb{R}} \mathcal{STF}_{\phi}^{\vartheta}\{h\}(t, \psi) K_{-\vartheta}(x, \psi) dx.$$

Thus, the desired result is obtained. □

Next, we will build Sharp Housdorff-Young inequality for the short-time fractional Fourier transform (STFrFT), which generalizes Sharp Hausdorff-Young inequality for the FT in StFrFT domains.

**Theorem 3.5.** *Let  $p(1 \leq r \leq 2)$  and  $q$  such that  $\frac{1}{p} + \frac{1}{q} = 1$ , then for any  $h, \phi \in L^p(\mathbb{R})$ , we have:*

$$\left\| \mathcal{STF}_\phi^\vartheta \{h\}(t, \psi) \right\|_{L^q(\mathbb{R} \times \mathbb{R})} \leq \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi} q^{\frac{1}{2q}}} \|h\|_{L^p(\mathbb{R})} \|\phi\|_{L^p(\mathbb{R})}. \quad (3.18)$$

**Proof.** With aid of Sharp Housdorff-Young inequality for the FT, we find that:

$$\left( \int_{\mathbb{R}} |\mathcal{F}\{h(x)\}(\psi)|^q d\psi \right)^{\frac{1}{q}} \leq \frac{p^{\frac{1}{2p}}}{q^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |h(x)|^p dx \right)^{\frac{1}{p}}. \quad (3.19)$$

We replace  $h(x)$  by  $\check{A}_{h,\phi}^\vartheta(x, t)$  by (3.6) into both sides of equation (3.19) above, we get:

$$\left( \int_{\mathbb{R}} |\mathcal{F}\{\check{A}_{h,\phi}^\vartheta\}(\psi)|^q d\psi \right)^{\frac{1}{q}} \leq \frac{p^{\frac{1}{2p}}}{q^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |\check{A}_{h,\phi}^\vartheta(x, t)|^p dx \right)^{\frac{1}{p}}. \quad (3.20)$$

We set  $\psi = \psi \csc \vartheta$  in equation (3.20) above, we obtain that:

$$\left( \int_{\mathbb{R}} |\mathcal{F}\{\check{A}_{h,\phi}^\vartheta\}(\psi \csc \vartheta)|^q d(\psi \csc \vartheta) \right)^{\frac{1}{q}} \leq \frac{p^{\frac{1}{2p}}}{q^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |\check{A}_{h,\phi}^\vartheta(x, t)|^p dx \right)^{\frac{1}{p}}. \quad (3.21)$$

Applying (3.5) to left hand side of equation (3.21), we find:

$$\left( (2\pi)^{\frac{q}{2}} |\sin \vartheta|^{\frac{q}{2}-1} \int_{\mathbb{R}} \left| \mathcal{STF}_\phi^\vartheta \{h\}(t, \psi) \right|^q d\psi \right)^{\frac{1}{q}} \leq \frac{p^{\frac{1}{2p}}}{q^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |h(x)|^p |\overline{\phi(x-t)}|^p dx \right)^{\frac{1}{p}}. \quad (3.22)$$

Hence,

$$\left( \int_{\mathbb{R}} \left| \mathcal{STF}_\phi^\vartheta \{h\}(t, \psi) \right|^q d\psi \right)^{\frac{1}{q}} \leq \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi} q^{\frac{1}{2q}}} \left( \int_{\mathbb{R}} |h(x)|^p |\overline{\phi(x-t)}|^p dx \right)^{\frac{1}{p}}, \quad (3.23)$$

and

$$\int_{\mathbb{R}} \left| \mathcal{STF}_\phi^\vartheta \{h\}(t, \psi) \right|^q d\psi \leq \left( \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi} q^{\frac{1}{2q}}} \right)^q \left( \int_{\mathbb{R}} |h(x)|^p |\overline{\phi(x-t)}|^p dx \right)^{\frac{q}{p}}. \quad (3.24)$$

Integrating both sides equation (3.24) with respect to  $dt$  we get:

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STF}_\phi^\vartheta \{h\}(t, \psi) \right|^q d\psi dt \leq \left( \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi} q^{\frac{1}{2q}}} \right)^q \left( \int_{\mathbb{R}} \int_{\mathbb{R}} |h(x)|^p |\overline{\phi(x-t)}|^p dx dt \right)^{\frac{q}{p}}. \quad (3.25)$$

Equation (3.25) may be expressed as:

$$\begin{aligned} \left( \int_{\mathbb{R}} \int_{\mathbb{R}} \left| \mathcal{STFr}_{\phi}^{\vartheta} \{h\}(t, \psi) \right|^q d\psi dt \right)^{\frac{1}{q}} &\leq \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi q^{\frac{1}{2q}}}} \left( \int_{\mathbb{R}} |h(x)|^p dx \int_{\mathbb{R}} |\overline{\phi(x-t)}|^p dt \right)^{\frac{1}{p}} \\ &= \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi q^{\frac{1}{2q}}}} \|f\|_{L^p(\mathbb{R})} \|\phi\|_{L^p(\mathbb{R})}. \end{aligned} \quad (3.26)$$

The above equation is the same as:

$$\left\| \mathcal{STFr}_{\phi}^{\vartheta} \{h\}(t, \psi) \right\|_{L^q(\mathbb{R} \times \mathbb{R})} \leq \frac{|\sin \vartheta|^{\frac{1}{q} - \frac{1}{2}} p^{\frac{1}{2p}}}{\sqrt{2\pi q^{\frac{1}{2q}}}} \|h\|_{L^p(\mathbb{R})} \|\phi\|_{L^p(\mathbb{R})}.$$

We obtained the desired result.  $\square$

#### 4. Conclusion

In this paper, we have introduced the short-time fractional Fourier transform (STFrFT) and investigated its essential properties. We have investigated Sharp Housdorff-Young inequality related to this transformation by applying several relations and properties that have been obtained for the proposed transformation.

#### 5. Acknowledgments

The authors are thankful to St. Nurhilmah Busrah for assistance in writing the manuscript.

#### Bibliography

- [1] Namias, V., 1980, The fractional order Fourier transform and its application to quantum mechanics, *J. Inst. Math. Appl.*, Vol. **25**: 241–265
- [2] McBride, A.C., Kerr, F.H., 1987, On Namias's fractional Fourier transforms, *IMA J. Appl. Math.*, Vol **39**: 159 – 175
- [3] Tao, R., Xin, Y., and Wang, Y., 2007, Double image encryption based on random phase encoding in the fractional Fourier domain, *Opt. Exp.*, Vol. **15**: 16067 - 16079
- [4] Tao, R., Lang, J., and Wang, Y., 2008, Optical image encryption based on the multiple-parameter fractional Fourier transform, *Opt. Lett.*, Vol. **33**: 581 - 583
- [5] Qi, L., Tao, R., Zhou, S.Y., and Wang, Y., 2004, Detection and parameter estimation of multicomponent LFM signal based on the fractional Fourier transform, *Science in China (Ser. F. Information Sciences)*, Vol. **47**: 184 - 198
- [6] Pei, S.C., and Ding, J.J., 2000, Closed-form discrete fractional and affine Fourier transforms, *IEEE Transactions on Signal Processing*, Vol. **48**(5): 1338 - 1353
- [7] Durak, L., and Arikan, O., 2003, Short-time Fourier transform: Two fundamental properties and an optimal implementation, *IEEE Transactions on Signal Processing*, Vol. **51**(5): 1231 - 1242
- [8] Bahri, M., 2017, Logarithmic uncertainty principle, convolution theorem related to continuous fractional wavelet transform and its properties on a generalized Sobolev space, *International Journal of Wavelets, Multiresolution and Information Processing*, Vol. **15**(5): 1750050

- [9] Anh, P.K., Castro, L.P., Thao, P.T., 2017, Two new convolutions for the fractional Fourier transform, *Wireless Personal Communications*, Vol. **92**: 623 – 637
- [10] Shi, J., Sha, X., Shong, X., and Zhang, N., 2012, Generalized convolution theorem associated with fractional Fourier transform, *Wireless Communication and Mobile Computing*, Vol. **14**: 1340 – 1351
- [11] Tao, R., Li, Y. L., Wang, Y., 2009, Short-time fractional Fourier transform and its applications, *IEEE Transactions on Signal Processing*, Vol. **58**(5): 2568 – 2580
- [12] Tao, R., Qi L., and Wang, Y. 2004, *Theory and Application of the Fractional Fourier Transform*, Tsinghua University Press, Beijing.
- [13] Cohen, L., 2000, The uncertainty principles of windowed wave functions, *Opt. Communication*, **179**(2000): 221 - 229
- [14] Wódkiewicz, K., 1984, Operational approach to phase-space measurements in quantum mechanics, *Phys. Rev. Lett.*, Vol. **52**(13): 1064 - 1067
- [15] Stankovic, L., Alieva, T., Bastiaans, M.J., 2003, Timefrequency signal analysis based on the windowed fractional Fourier transform, *Signal Processing*, Vol. **8**: 2459 - 2468
- [16] Guanlei, X., Xiaotong, W., Xiaogang, X., 2009, The logarithmic, Heisenberg's and short-time uncertainty principles associated with fractional Fourier transform, *Signal Processing*, Vol. **89**(3): 339 – 343
- [17] Bahri, M., and Karim, S.A.A., 2023, Fractional Fourier transform: main properties and inequalities, *Mathematics*, Vol. **11**(5): 1234